Author Index

AKASHI T., NAGAFUCHI S., ANZAI K., KITAMURA D., WANG J., TANIUCHI I., NIHO Y. & WATANABE T. Proliferation of $CD3^+$ $B220^-$ single-positive normal T cells was suppressed in B-cell-deficient *lpr* mice, 238
ALEXANDER J. *see* BREWER J.M.
ANZAI K. *see* AKASHI T.
AOKI Y. *see* YAMAZAKI T.
APPARAILLY F. *see* JORGENSEN C.
ARAGANE Y. *see* MÖLLER A.
ARIZONO N. *see* MORIMOTO M.
ASANO T. *see* SHIBATA S.
ASEA A. & STEIN-STREILEIN J. Signalling through NK1.1 triggers NK cells to die but induces NK T cells to produce interleukin-4, 296
ASHERSON G.L. *see* DIELI F.
ATEN J. *see* ROOS A.

BADMINTON M.N. *see* SALA-NEWBY G.B.
BARATA L.T. *see* YING S.
BARKANS J. *see* YING S.
BARTELS T., STEINBACH F., HAHN G., LUDWIG H. & BORCHERS K. *In situ* study on the pathogenesis and immune reaction of equine herpesvirus type 1 (EHV-1) infections in mice, 329
BECKER D. *see* SALOGA J.
BEECH J.T. *see* SIEW L.K.
BEFUS A.D. *see* MACDONALD A.J.
BELL E.B. *see* SPARSHOTT S.M.
BELLINGHAUSEN I. *see* SALOGA J.
BEN-MOSHE O. *see* EREN R.
BETTS C. *see* COULSON P.S.
BETTS C.J. & WILSON R.A. Th1 cytokine mRNA expression dominates in the skin-draining lymph nodes of C57BL/6 mice following vaccination with irradiated *Schistosoma mansoni* cercariae, but is down-regulated upon challenge infection, 49
BISSONNETTE E.Y. *see* MACDONALD A.J.
BLAKEMORE W.F. *see* BORROW P.
BLUETHMANN H. *see* BREWER J.M.
BORCHERS K. *see* BARTELS T.
BORLAND G., ROSS J.A. & GUY K. Forms and functions of CD44, 139
BORROW P., WELSH C.J.R., TONKS P., DEAN D., BLAKEMORE W.F. & NASH A.A. Investigation of the role of delayed-type-hypersensitivity responses to myelin in the pathogenesis of Theiler's virus-induced demyelinating disease, 478
BREIT T.M. *see* VERSCHUREN M.C.M.
BREWER J.M., CONACHER M., GAFFNEY M., DOUGLAS M., BLUETHMANN H. & ALEXANDER J. Neither interleukin-6 nor signalling via tumour necrosis factor receptor-1 contribute to the adjuvant activity of Alum and Freund's adjuvant, 41
BRIEN J.H. *see* SHIRZADEH H.
BROOK M.A. *see* HERITAGE P.L.
BUISMAN A.M., LANGERMANS J.A.M. & VAN FURTH R. Effect of granulocyte–macrophage colony-stimulating factor on the number of leucocytes and course of *Listeria monocytogenes* infection in naive and leucocytopenic mice, 73
BULMER J.N. *see* JAKOBSEN P.H.
BUNYAN K. *see* HAIG D.M.
BURTON R.C. *see* SHIRZADEH H.

CAMPBELL A.K. *see* SALA-NEWBY G.B.
CANOVAS F. *see* JORGENSEN C.
CAO Y. *see* YANG M.
CAPUTI A.P. *see* CUZZUCREA S.
CARNAHAN J.A. *see* VARADE W.S.
CHAND M.A. *see* ROOS A.
CHEEVER M.A. *see* DISIS M.L.
CHEN Z. *see* GORCZYNSKI R.M.
CHOW S.C. *see* OSORIO L.M.
CHUA-INTRA B., IVANYI J., HILLS A., THOLE J., MORENO C. & VORDERMEIER H. M. Predominant recognition of species-specific determinants of the GroES homologues from *Mycobacterium leprae* and *M. tuberculosis*, 64
CIRCOLO A. *see* WANG X.
CLAESSEN N. *see* ROOS A.
CLARK M.R. *see* REDPATH S.
COLTEN H.R. *see* WANG X.
CONACHER M. *see* BREWER J.M.
COPIER J., POTTER P., SACKS S.H. & KELLY A.P. Multiple signals regulate the intracellular trafficking of HLA-DM in B-lymphoblastoid cells, 505
CORNACCHIONE P., SCARINGI L., FETTUCCIARI K., ROSATI E., SABATINI R., OREFICI G., VON HUNOLSTEIN C., MODESTI A., MODICA A., MINELLI F. & MARCONI P. Group B streptococci persist inside macrophages, 86
COULSON P.S., SMYTHIES L.E., BETTS C., MABBOTT N.A., STERNBERG J.M., WEI X.-G., LIEW F.Y. & WILSON R. A. Nitric oxide produced in the lungs of mice immunized with the radiation-attenuated schistosome vaccine is not the major agent causing challenge parasite elimination, 55
COURET I. *see* JORGENSEN C.
COWARD A. *see* GUGASYAN R.
CUZZUCREA S., CAPUTI A.P. & ZINGARELLI B. Peroxynitrite-mediated DNA strand breakage activates poly (ADP-ribose) synthetase and causes cellular energy depletion in carrageenan-induced pleurisy, 96

DAGAN S. *see* EREN R.
DALGAARD T.S. *see* LAURSEN S.B.
DAMGAARD G. *see* OLESEN E.H.
DAUDI N. *see* EREN R.
DAY N.K. *see* KIZAKI T.
DE GREEF C. *see* ZHU D.
DE SILVA H.D., VAN DRIEL I.R., LA GRUTA N., TOH B.H. & GLEESON P.A. $CD4^+$ T cells, but not $CD8^+$ T cells, are required for the development of experimental autoimmune gastritis, 405
DEAN D. *see* BORROW P.
DIELI F., PTAK W., SIRECI G., ROMANO G.C., POTESTIO M., SALERNO A. & ASHERSON G.L. Cross-talk between $V\beta8^+$ and $\gamma\delta^+$ T lymphocytes in contact sensitivity, 469
DISIS M.L., SHIOTA F.M. & CHEEVER M. A. Human HER-2/neu protein immunization circumvents tolerance to rat neu: a vaccine strategy for 'self' tumour antigens, 192
DOEBERITZ M.V.K. SEE LINDAUER M.
DOI K. *see* SHIBATA S.
DONACHIE A.M. *see* SMITH R.E.
DOUGLAS M. *see* BREWER J.M.
DOYLE A.G., RAMM L. & KELSO A. The $CD4^+$ T-cell response to protein immunization is independent of accompanying IFN-γ-producing $CD8^+$ T cells, 341
DURHAM S.R. *see* YING S.

EASTCOTT J.W. *see* KAWAI T.
EHRLICH S. *see* VON BONIN A.
ELSON C.J. *see* SIEW L.K.
ENK A.H. *see* SALOGA J.

EREN R., LUBIN I., TERKIELTAUB D., BEN-MOSHE O., ZAUBERMAN A., UHLMANN R., TZAHOR T., MOSS S., ILAN E., SHOUVAL D., GALUN E., DAUDI N., MARCUS H., REISNER Y. & DAGAN S. Human monoclonal antibodies specific to hepatitis B virus generated in a human/mouse radiation chimera: the Trimera system, 154

FERRANTE P. *see* SPECIALE L.
FETTUCCIARI K. *see* CORNACCHIONE P.
FILSINGER S. *see* KRAFT M.
FLEISCHER B. *see* VON BONIN A.
FROELICH C.J. *see* WEVER P.C.
FROLOV I., HOURI-HADAD Y., SOSKOLNE A. & SHAPIRA L. *In vivo* exposure to *Porphyromonas gingivalis* up-regulates nitric oxide but suppresses tumour necrosis factor-α production by cultured macrophages, 323
FUJIMOTO J. *see* IWASAKI T.

GAFFNEY M. *see* BREWER J.M.
GALUN E. *see* EREN R.
GARDINER C.M. & REEN D.J. Differential cytokine regulation of natural killer cell-mediated necrotic and apoptotic cytotoxicity, 511
GARZA A. *see* LIANG B.
GIRVIN R.M. *see* LIGHTBODY K.A.
GLEESON P.A. *see* DE SILVA H.D.
GONZÁLEZ-FERNÁNDEZ A. & MILSTEIN C. Low antigen dose favours selection of somatic mutants with hallmarks of antibody affinity maturation, 149
GOOD R.A. *see* KIZAKI T.
GORCZYNSKI L. *see* GORCZYNSKI R.M.
GORCZYNSKI R.M., CHEN Z., ZENG H., GORCZYNSKI L. & TERZIOGLU E. Analysis of cytokine production and Vβ T-cell receptor subsets in irradiated recipients receiving portal or peripheral venous reconstitution with allogeneic bone marrow cells, with or without additional anti-cytokine monoclonal antibodies, 221
GOTHEFORS L. *see* RYAN M.
GOYAL P.K. *see* ISHIKAWA N.
GRANT J.A. *see* YING S.
GREEN C.B. *see* WHALEN M.M.
GREENWOOD B.M. *see* JAKOBSEN P.H.
GRÜTZKAU A. *see* MÖLLER A.
GÜCKEL B. *see* LINDAUER M.
GUGASYAN R., COWARD A., O'CONNOR L., SHORTMAN K. & SCOLLAY R. Emigration of mature T cells from the thymus is inhibited by the imidazole-based compound 2-acetyl-4 tetrahydroxybutylimidazole, 398
GUY K. *see* BORLAND G.

HACK C.E. *see* WEVER P.C.
HAHN G. *see* BARTELS T.
HAIG D.M., McINNES C.J., THOMSON J., WOOD A., BUNYAN K. & MERCER A. The orf virus OV20.0L gene product is involved in interferon resistance and inhibits an interferon-inducible, double-stranded RNA-dependent kinase, 335
HAMADA J.-I. *see* LI Y.-Q.
HAMANA T. *see* LAURSEN S.B.
HAMANO T. *see* IWASAKI T.
HAMEL M.E., NOTEBOOM E. & KRUISBEEK A. M. Non-responsiveness of antigen-experienced CD4 T cells reflects more stringent co-stimulatory requirements, 366
HAMMARSTRÖM L. *see* MOUSAVI M.
HÄNSCH G.M. *see* KRAFT M.
HARA T., SUZUKI Y., NAKAZAWA S., NISHIMURA H., NAGASAWA S., NISHIGUCHI M., MATSUMOTO M., HATANAKA M., KITAMURA M. & SEYA T. Post-translational modification and intracellular localization of a splice product of CD46 cloned from human testis: role of the intracellular domains in O-glycosylation, 546
HATANAKA M. *see* HARA T.
HAYASHI T. *see* MORIMOTO M.
HENZ B.M. *see* MÖLLER A.
HERITAGE P.L., BROOK M.A., UNDERDOWN B.J. & McDERMOTT M. R. Intranasal immunization with polymer-grafted microparticles activates the nasal-associated lymphoid tissue and draining lymph nodes, 249
HILLS A. *see* CHUA-INTRA B.
HIRANO S. Nitric oxide-mediated cytotoxic effects of alveolar macrophages on transformed lung epithelial cells are independent of the β_2 integrin-mediated intercellular adhesion, 102
HIRATA M. *see* ITO H.-O.
HOLMES C.H. *see* WAINWRIGHT S.D.
HOLMSKOV U., JENSENIUS J.C., TORNØE I. & LØVENDAHL P. The plasma levels of conglutinin are heritable in cattle and low levels predispose to infection, 431
HOSOKAWA M *see* LI Y.-Q.
HOURI-HADAD Y. *see* FROLOV I.

ILAN E. *see* EREN R.
IMOTO T. *see* ITO H.-O.
INSEL R.A. *see* VARADE W.S.
INSERRA P. *see* LIANG B.
ISHIKAWA N., GOYAL P.K., MAHIDA Y.R., LI K.-F. & WAKELIN D. Early cytokine responses during intestinal parasitic infections, 257
ITO H.-O., SO T., HIRATA M., KOGA T., UEDA T. & IMOTO T. Tolerogenic activity of polyethylene glycol-conjugated lysozyme distinct from that of the native counterpart, 200
ITOH Y. *see* KIZAKI T.
IVANYI J. *see* CHUA-INTRA B.
IWABUCHI K. *see* KIZAKI T.
IWASAKI T., HAMANO T., FUJIMOTO J. & KAKISHITA E. Regulation of interleukin-6 and interleukin-6R α (gp80) expression by murine immunoglobulin-secreting B-cell hybridomas, 498

JAKOBSEN P.H., RASHEED F.N., BULMER J.N., THEISEN M., RIDLEY R.G. & GREENWOOD B. M. Inflammatory reactions in placental blood of *Plasmodium falciparum*-infected women and high concentrations of soluble E-selectin and a circulating *P. falciparum* protein in the cord sera, 264
JAQUET C. *see* JORGENSEN C.
JENSEN T.V. *see* LAURSEN S.B.
JENSENIUS J C. *see* HOLMSKOV U.
see also LAURSEN S.B.
JIANG S. *see* LIANG B.
JOHNSON A.A. *see* OLESEN E.H.
JONDAL M. *see* OSORIO L.M.
JORGENSEN C., APPARAILLY F., COURET I., CANOVAS F., JAQUET C. & SANY J. Interleukin-4 and interleukin-10 are chondroprotective and decrease mononuclear cell recruitment in human rheumatoid synovium *in vitro*, 518
JULKUNEN I. *see* SARENEVA T.
JUUL-MADSEN H.R. *see* LAURSEN S.B.

KABELITZ D. *see* KRAFT M.
KAKISHITA E. *see* IWASAKI T.
KAWAI T., SHIMAUCHI H., EASTCOTT J.W., SMITH D.J. & TAUBMAN M. A. Antigen direction of specific T-cell clones into gingival tissues, 11
KAWAKAMI M. *see* LAURSEN S.B.
KAY A.B. *see* YING S.
KELLY A.P. *see* COPIER J.
KELSO A. *see* DOYLE A.G.
KIM H.-M., KIM K.-S. & LEE E.-H. Specific inhibition of immunoglobin E-mediated allergic reaction using antisense FcεRIα oligodeoxynucleotides, 589
KIM K.-S. *see* KIM H.-M.
KIMURA K. *see* LI Y.-Q.
KING C.A. *see* ZHU D.
KINGSLEY P.D. *see* VARADE W.S.
KITAMURA D. *see* AKASHI T.
KITAMURA M. *see* HARA T.
KIZAKI T., OOKAWARA T., OH-ISHI S., ITOH Y., IWABUCHI K., ONOÉ K., DAY N.K., GOOD R.A. & OHNO H. An increase in basal glucocorticoid concentration with age induces suppressor macrophages with high-density FcγRII/III, 409
KNOP J. *see* SALOGA J.

KOBAYASHI M. *see* LI Y.-Q.
KOGA T. *see* ITO H.-O.
KRAFT M., FILSINGER S., KRÄMER K.-L., KABELITZ D., HÄNSCH G.M. & SCHOELS M. Synovial fibroblasts as target cells for staphylococcal enterotoxin-induced T-cell cytotoxicity, 20
KRÄMER K.-L. *see* KRAFT M.
KRÜGER-KRASAGAKES S. *see* MÖLLER A.
KRUISBEEK A.M. *see* HAMEL M.E.
KÜHN S. *see* SALOGA J.

LA GRUTA N. *see* DE SILVA H.D.
LAKKIS F.G. *see* ROOS A.
LANGERMANS J.A.M. *see* BUISMAN A.M.
LAURSEN S.B., DALGAARD T.S., THIEL S., LIM B.-L., JENSEN T.V., JUUL-MADSEN H.R., TAKAHASHI A., HAMANA T., KAWAKAMI M. & JENSENIUS J. C. Cloning and sequencing of a cDNA encoding chicken mannan-binding lectin (MBL) and comparison with mammalian analogues, 421
LEE E.-H. *see* KIM H.-M.
LEE J. *see* LIANG B.
LESLIE R.G.Q. *see* OLESEN E.H.
LI K.-F. *see* ISHIKAWA N.
LI Y.-Q., KOBAYASHI M., YUAN L., WANG J., MATSUSHITA K., HAMADA J.-I., KIMURA K., YAGITA H., OKUMURA K. & HOSOKAWA M. Protein kinase C mediates the signal for interferon-γ mRNA expression in cytotoxic T cells after their adhesion to laminin, 455
LIANG B., ZHANG Z., INSERRA P., JIANG S., LEE J., GARZA A., MARCHALONIS J.J. & WATSON R.R. Injection of T-cell receptor peptide reduces immunosenescence in aged C57BL/6 mice, 462
LIEW F.Y *see* COULSON P.S.
LIGHTBODY K.A., GIRVIN R.M., POLLOCK D.A., MACKIE D.P., NEILL S.D. & POLLOCK J. M. Recognition of a common mycobacterial T-cell epitope in MPB59 of *Mycobacterium bovis*, 314
LIM B.-L. *see* LAURSEN S.B.
LINDAUER M., RUDY W., GÜCKEL B., DOEBERITZ M.V.K., MEUER S.C. & MOEBIUS U. Gene transfer of costimulatory molecules into a human colorectal cancer cell line: requirement of CD54, CD80 and class II MHC expression for enhanced immunogenicity, 390
LIPPERT U. *see* MÖLLER A.
LOKKI M.-L. *see* WANG X.
LØVENDAHL P. *see* HOLMSKOV U.
LUBIN I. *see* EREN R.
LUDWIG H. *see* BARTELS T.

MABBOTT N.A. *see* COULSON P.S.
MACDONALD A.J., PICK J., BISSONNETTE E.Y. & BEFUS A.D. Rat mucosal mast cells: the cultured bone marrow-derived mast cell is biochemically and functionally analogous to its counterpart *in vivo*, 533
MACKIE D.P. *see* LIGHTBODY K.A.
MAHIDA Y.R. *see* ISHIKAWA N.
MARCHALONIS J.J. *see* LIANG B.
MARCONI P. *see* CORNACCHIONE P.
MARCUS H. *see* EREN R.
MATIKAINEN S. *see* SARENEVA T.
MATSUMOTO M. *see* HARA T.
MATSUSHITA K. *see* LI Y.-Q.
McDERMOTT M.R. *see* HERITAGE P.L.
McINNES C.J. *see* HAIG D.M.
McLAREN F.H. *see* SMITH R.E.
MEDINA E. & NORTH R. J. Resistance ranking of some common inbred mouse strains to *Mycobacterium tuberculosis* and relationship to major histocompatability complex haplotype and *Nramp1* genotype, 270
MELÉN K. *see* SARENEVA T.
MENG Q. *see* YING S.
MERCER A. *see* HAIG D.M.
MEUER S.C. *see* LINDAUER M.
MICHAELSEN T. *see* REDPATH S.
MILLER E. *see* RYAN M.
MILLS K.H.G. *see* RYAN M.
MILSTEIN C. *see* GONZÁLEZ-FERNÁNDEZ A.
MINELLI F. *see* CORNACCHIONE P.
MODESTI A. *see* CORNACCHIONE P.
MODICA A. *see* CORNACCHIONE P.
MOEBIUS U. *see* LINDAUER M.
MOHAPATRA S.S. *see* YANG M.
MÖLLER A., HENZ B.M., GRÜTZKAU A., LIPPERT U., ARAGANE Y., SCHWARZ T. & KRÜGER-KRASAGAKES S. Comparative cytokine gene expression: regulation and release by human mast cells, 289
MORENO C. *see* CHUA-INTRA B.
MORIMOTO M., YAMADA M., ARIZONO N. & HAYASHI T. Lactic dehydrogenase virus infection enhances parasite egg production and inhibits eosinophil and mast cell responses in mice infected with the nematode *Nippostrongylus brasiliensis*, 540
MOSS S. *see* EREN R.
MOUSAVI M., RABBANI H., PILSTRÖM L. & HAMMARSTRÖM L. Characterization of the gene for the membrane and secretory form of the IgM heavy-chain constant region gene (Cμ) of the cow (*Bos taurus*), 581
MOWAT A.MCI. *see* SMITH R.E.
MUHUMUZA L., SEGRE D. & SEGRE M. Antibodies with idiotypic and anti-idiotypic reactivity (epibodies) in conventional immune responses to dinitrophenylated carriers, 572
MURPHY G. *see* RYAN M.

NAGAFUCHI S. *see* AKASHI T.
NAGASAWA S. *see* HARA T.
NAITO M. *see* SHIBATA S.
NAKAZAWA S. *see* HARA T.
NASH A.A. *see* BORROW P.
NATVIG J.B. *see* THORPE S.J.
NEILL S.D. *see* LIGHTBODY K.A.
NIHO Y. *see* AKASHI T.
NILSSON I.-M. *see* ZHAO Y.-X.
NILSSON L. *see* RYAN M.
NISHIGUCHI M. *see* HARA T.
NISHIMURA H. *see* HARA T.
NOGUCHI A. *see* SHIBATA S.
NORTH R.J. *see* MEDINA E.
NOTEBOOM E. *see* HAMEL M.E.

O'CONNOR L. *see* GUGASYAN R.
O'HAGAN D.T. *see* UGOZZOLI M.
OGURA A. *see* SHIBATA S.
OH-ISHI S. *see* KIZAKI T.
OHNO H. *see* KIZAKI T.
OKAMOTO Y. *see* TAKEICHI O.
OKUMURA K. *see* LI Y.-Q.
OLESEN E.H., JOHNSON A.A., DAMGAARD G. & LESLIE R. G. Q. The requirement of localized, CR2-mediated, alternative pathway activation of complement for covalent disposition of C3 fragments on normal B cells, 177
ONOÉ K. *see* KIZAKI T.
OOKAWARA T. *see* KIZAKI T.
OREFICI G. *see* CORNACCHIONE P.
OSORIO L.M., ROTTENBERG M., JONDAL M. & CHOW S. C. Simultaneous cross-linking of CD6 and CD28 induces cell proliferation in resting T cells, 358
OTT G.S. *see* UGOZZOLI M.
ØYMAR K. *see* RYAN M.

PARKHOUSE R.M.E. *see* YANG H.
PASCUAL D.W. *see* ROOS A.
PELKONEN J. *see* SARENEVA T.
PICK J. *see* MACDONALD A.J.
PILSTRÖM L. *see* MOUSAVI M.
POLLOCK D.A. *see* LIGHTBODY K.A.
POLLOCK J.M. *see* LIGHTBODY K.A.
POTESTIO M. *see* DIELI F.
POTTER P. *see* COPIER J.
PTAK W. *see* DIELI F.

RABBANI H. *see* MOUSAVI M.
RADL J. *see* ZHU D.
RAMM L. *see* DOYLE A.G.
RASHEED F.N. *see* JAKOBSEN P.H.
REDPATH S., MICHAELSEN T., SANDLIE I. & CLARK M.R. Activation of complement by human IgG1 and human IgG3 antibodies against the human leucocyte antigen CD52, 595
REEN D.J. *see* GARDINER C.M.
REISNER Y. *see* EREN R.
REMBOLD C.M. *see* SALA-NEWBY G.B.
RIDLEY R.G. *see* JAKOBSEN P.H.
RODA K. *see* SPECIALE L.
ROMANO G.C. *see* DIELI F.
ROOK G.A.W. *see* WANG C.-C.
ROOS A., SCHILDER-TOL E.J.M., CHAND M.A., CLAESSEN N., LAKKIS F.G., PASCUAL D.W., WEENING J.J. & ATEN J. Differential regulation of expression of the MHC class II molecules RT1.B and RT1.D on rat B lymphocytes: effects of interleukin-4, interleukin-13 and interferon-γ, 33
ROSATI E. *see* CORNACCHIONE P.
ROSS J.A. *see* BORLAND G.
ROTTENBERG M. *see* OSORIO L.M.
RUDY W. *see* LINDAUER M.
RYAN E. *see* RYAN M.
RYAN M., MURPHY G., RYAN E., NILSSON L., SHACKLEY F., GOTHEFORS L., ØYMAR K., MILLER E., STORSAETER J. & MILLS K. H. G. Distinct T-cell subtypes induced with whole cell and acellular pertussis vaccines in children, 1

SABATINI R. *see* CORNACCHIONE P.
SACKS S.H. *see* COPIER J.
SAITO I. *see* TAKEICHI O.
SAITO T. *see* TAKEICHI O.
SALA-NEWBY G.B., TAYLOR K.M., BADMINTON M.N., REMBOLD C.M. & CAMPBELL A.K. Imaging bioluminescent indicators shows Ca^{2+} and ATP permeability thresholds in live cells attacked by complement, 601
SALERNO A. *see* DIELI F.
SALOGA J., ENK A.H., BECKER D., BELLINGHAUSEN I., KÜHN S. & KNOP J. Active suppression induced by cutaneous exposure to bacterial superantigen is prevented by interleukin-12 treatment *in vivo*, 485
SANDLIE I. *see* REDPATH S.
SANY J. *see* JORGENSEN C.
SARENEVA T., MATIKAINEN S., VANHATALO J., MELÉN K., PELKONEN J. & JULKUNEN I. Kinetics of cytokine and NFAT gene expression in human interleukin-2-dependent T lymphoblasts stimulated via T-cell receptor, 350
SARESELLA M. *see* SPECIALE L.
SCARINGI L. *see* CORNACCHIONE P.
SCHILDER-TOL E.J.M. *see* ROOS A.
SCHOELS M. *see* KRAFT M.
SCHWARZ T. *see* MÖLLER A.
SCOLLAY R. *see* GUGASYAN R.
SEGRE D. *see* MUHUMUZA L.
SEGRE M. *see* MUHUMUZA L.
SEYA T. *see* HARA T.
SHACKELFORD P.G. *see* WANG X.
SHACKLEY F. *see* RYAN M.
SHAPIRA L. *see* FROLOV I.
SHIBATA S., ASANO T., NOGUCHI A., NAITO M., OGURA A. & DOI K. Peritoneal macrophages play an important role in eliminating human cells from severe combined immunodeficient mice transplanted with human peripheral blood lymphocytes, 524
SHIMAUCHI H. *see* KAWAI T.
SHIOTA F.M. *see* DISIS M.L.
SHIRZADEH H., BURTON R.C., BRIEN J.H. & SMART Y. C. Monoclonal antibody anti-NC-2 identifies a second receptor on cells mediating natural cytotoxicity in mice, 122
SHORTMAN K. *see* GUGASYAN R.
SHOUVAL D. *see* EREN R.
SIEW L.K., BEECH J.T., THOMPSON S.J. & ELSON C.J. Effect of T-helper cytokine environment on specificity of T-cell responses to mycobacterial 65 000 MW heat-shock protein, 493
SIMPSON K.L. *see* WAINWRIGHT S.D.
SIRECI G. *see* DIELI F.
SMART Y.C. *see* SHIRZADEH H.
SMITH D.J. *see* KAWAI T.
SMITH R.E., DONACHIE A.M., McLAREN F.H. & MOWAT A.MCI. Preservation of mucosal and systemic adjuvant properties of ISCOMS in the absence of functional interleukin-4 or interferon-γ, 556
SMYTHIES L.E. *see* COULSON P.S.
SO T. *see* ITO H.-O.
SOSKOLNE A. *see* FROLOV I.
SPAENY L.H.A. *see* WEVER P.C.
SPARSHOTT S.M. & BELL E.B. Lymphocyte trafficking: CD4 T cells with a 'memory' phenotype ($CD45RC^-$) freely cross lymph node high endothelial venules *in vivo*, 447
SPECIALE L., RODA K., SARESELLA M., TARAMELLI D. & FERRANTE P. Different endothelins stimulate cytokine production by peritoneal macrophages and microglial cell line, 109
SPELLERBERG M.B. *see* THORPE S.J.
STEINBACH F. *see* BARTELS T.
STEIN-STREILEIN J. *see* ASEA A.
STERNBERG J.M. *see* COULSON P.S.
STEVENSON F.K. *see* THORPE S.J.
see also ZHU D.
STORSAETER J. *see* RYAN M.
SUTTON K.A., VU M.N. & WILKINSON M. F. Distal Vβ promoters transcribe novel T-cell receptor-β transcripts in early development, 213
SUZUKI Y. *see* HARA T.

TAKAHASHI A. *see* LAURSEN S.B.
TAKEICHI O., SAITO I., OKAMOTO Y., TSURUMACHI T. & SAITO T. Cytokine regulation on the synthesis of nitric oxide *in vivo* by chronically infected human polymorphonuclear leucocytes, 275
TANIUCHI I. *see* AKASHI T.
TARAMELLI D. *see* SPECIALE L.
TARKOWSKI A. *see* ZHAO Y.-X.
TAUBMAN M.A. *see* KAWAI T.
TAYLOR K.M. *see* SALA-NEWBY G.B.
TEN BERGE I.J.M. *see* WEVER P.C.
TERKIELTAUB D. *see* EREN R.
TERZIOGLU E. *see* GORCZYNSKI R.M.
THEISEN M. *see* JAKOBSEN P.H.
THIEL S. *see* LAURSEN S.B.
THIELEMANS K. *see* ZHU D.
THOLE J. *see* CHUA-INTRA B.
THOMPSON K.M. *see* THORPE S.J.
THOMPSON S.J. *see* SIEW L.K.
THOMSON J. *see* HAIG D.M.
THORPE R. *see* THORPE S.J.
THORPE S.J., TURNER C.E., STEVENSON F.K., SPELLERBERG M.B., THORPE R., NATVIG J.B. & THOMPSON K. M. Human monoclonal antibodies encoded by the V4-34 gene segment show cold agglutinin activity and variable multireactivity which correlates with the predicted charge of the heavy-chain variable region, 129
TOH B.H. *see* DE SILVA H.D.
TONKS P. *see* BORROW P.
TORNØE I. *see* HOLMSKOV U.
TSURUMACHI T. *see* TAKEICHI O.
TURNER C.E. *see* THORPE S.J.
TZAHOR T. *see* EREN R.

UEDA T. *see* ITO H.-O.
UGOZZOLI M., O'HAGAN D.T. & OTT G.S. Intranasal immunization of mice with herpes simplex virus type 2 recombinant gD2: the effect of adjuvants on mucosal and serum antibody responses, 563
UHLMANN R. *see* EREN R.
UNDERDOWN B.J. *see* HERITAGE P.L.

VAN ARKEL C. *see* ZHU D.
VAN DER VLIET H.J.J. *see* WEVER P.C.
VAN DIEPEN F.N.J. *see* WEVER P.C.
VAN DONGEN J.J.M *see* VERSCHUREN M.C.M.
VAN DRIEL I.R. *see* DE SILVA H.D.
VAN FURTH R. *see* BUISMAN A.M.
VAN MEIRVENNE S. *see* ZHU D.
VANHATALO J. *see* SARENEVA T.
VARADE W.S., CARNAHAN J.A., KINGSLEY P.D. & INSEL R. A. Inherent properties of somatic hypermutation as revealed by human non-productive V_H6 immunoglobulin rearrangements, 171
VERSCHUREN M.C.M., WOLVERS-TETTERO I.L.M., BREIT T.M. & VAN DONGEN J.J.M. T-cell receptor Vδ-Jα rearrangements in human thymocytes: the role of Vδ-Jα rearrangements in T-cell receptor-δ gene deletion, 208
VON BONIN A., EHRLICH S. & FLEISCHER B. The transmembrane region of CD2-associated signal-transducing proteins is crucial for the outcome of CD2-mediated T-cell activation, 376
VON HUNOLSTEIN C. *see* CORNACCHIONE P.
VORDERMEIER H.M. *see* CHUA-INTRA B.
VU M.N. *see* SUTTON K.A.

WAINWRIGHT S.D., SIMPSON K.L. & HOLMES C. H. Calreticulin associates with non-HLA-A,-B class I proteins in the human choriocarcinoma cell lines JEG-3 and BeWo, 437
WAKELIN D. *see* ISHIKAWA N.
WANG C.-C. & ROOK G. A. W. Inhibition of an established allergic response to ovalbumin in BALB/c mice by killed *Mycobacterium vaccae*, 307
WANG J. *see* AKASHI T.
see also LI Y.-Q.
WANG X., CIRCOLO A., LOKKI M.-L., SHACKELFORD P.G., WETSEL R.A. & COLTEN H. R. Molecular heterogeneity in deficiency of complement protein C2 type I, 184
WATANABE T. *see* AKASHI T.
WATSON R.R. *see* LIANG B.
WEENING J.J. *see* ROOS A.
WEI X.-G. *see* COULSON P.S.
WELSH C.J.R. *see* BORROW P.
WETSEL R.A. *see* WANG X.
WEVER P.C., VAN DER VLIET H.J.J., SPAENY L.H.A., WOLBINK A.M., VAN DIEPEN F.N.J., FROELICH C.J., HACK C.E. & TEN BERGE I. J. M. The $CD8^+$ granzyme B^+ T-cell subset in peripheral blood from healthy individuals contains activated and apoptosis-prone cells, 383
WHALEN M.M. & GREEN C. B. Lysis-sensitive targets stimulate an elevation of cAMP in human natural killer cells, 415
WILKINSON M.F. *see* SUTTON K.A.
WILSON R.A. *see* BETTS C.J.
see also COULSON P.S.
WOLBINK A.M. *see* WEVER P.C.
WOLVERS-TETTERO I.L.M. *see* VERSCHUREN M.C.M.
WOOD A. *see* HAIG D.M.

YAGITA H. *see* LI Y.-Q.
YAMADA M. *see* MORIMOTO M.
YAMAZAKI T. & AOKI Y. Cathepsin G enhances human natural killer cytotoxicity, 115
YANG H. & PARKHOUSE R. M. E. Differential activation requirements associated with stimulation of T cells via different epitopes of CD3, 26
YANG M., CAO Y. & MOHAPATRA S.S. $CD8^+$ T cells inhibit immunoglobulin E synthesis in low responder SJL/J mice, 230
YING S., BARATA L.T., MENG Q., GRANT J.A., BARKANS J., DURHAM S.R. & KAY A. B. High-affinity immunoglobin E receptor ($Fc\varepsilon RI$)-bearing eosinophils, mast cells, macrophages and Langerhans' cells in allergen-induced late-phase cutaneous reactions in atopic subjects, 281
YUAN L. *see* LI Y.-Q.

ZAUBERMAN A. *see* EREN R.
ZENG H. *see* GORCZYNSKI R.M.
ZHANG Z. *see* LIANG B.
ZHAO Y.-X., NILSSON I.-M. & TARKOWSKI A. The dual role of interferon-γ in experimental *Staphylococcus aureus* septicemia versus arthritis, 80
ZHU D., VAN ARKEL C., KING C.A., VAN MEIRVENNE S., DE GREEF C., THIELEMANS K., RADL J. & STEVENSON F. K. Immunoglobulin V_H gene sequence analysis of spontaneous murine immunoglobulin-secreting B-cell tumours with clinical features of human disease, 162
ZINGARELLI B. *see* CUZZUCREA S.

Subject Index

Adjuvant activity, of Alum and Freund's:
role of interleukin-6 and tumour necrosis factor receptor-1, 41
Allergic reaction, immunoglobulin E mediated:
inhibition by antisense FcεRIα oligodeoxynucleotides, 589
Antibody affinity maturation:
selection of somatic mutants by low-dose antigen, 149
Autoimmune gastritis, experimental:
role of T cells, 405

B-cell-deficient *lpr* mice, 238

C2 deficiency:
type Ib, 184
Calreticulin:
association with non-HLA-A,-B class I proteins, 437
CD2-mediated T-cell activation, 376
CD3-induced T-cell activation, 26
CD44:
forms and functions, 139
CD46 splice product:
from human testis, 546
Complement activation, alternative pathway:
C3 deposition on B cells, 177
Complement activation by human IgG1 and IgG3 antibodies, 595
Complement attack:
permeability thresholds for Ca^{2+} and ATP, 601
Conglutinin, in cattle:
plasma levels, 431
Co-stimulatory molecules:
gene transfer, 390
Cytokine gene expression:
in T lymphocytes, 350
Cytokine induction:
by endothelins, 109
Cytokine mRNA expression in skin-draining lymph nodes:
after vaccination with irradiated *Schistosoma mansoni* cercariae, 49
Cytokine production and Vβ T-cell receptor subsets:
in bone marrow recipients, 221
Cytokine regulation:
of lymphokine-activated killer cell cytotoxicity, 511
Cytokine responses, early:
during intestinal parasitic infections, 257
Cytokine responses, T helper:
independent of $CD8^+$ cells, 341
Cytokine therapy:
in human rheumatoid synovium in severe combined immunodeficiency mice, 518

Delayed-type-hypersensitivity responses to myelin:
role in Theiler's disease, 478

Emigration, of mature T cells from the thymus:
inhibition by imidazole compound, 398
Epibodies:
in responses to dinitrophenylated carriers, 572
Equine herpesvirus type 1:
infection in mice, 329

FcεRI-bearing cells:
in allergen-induced late-phase response, 281

Granulocyte–macrophage colony-stimulating factor:
effect on *Listeria monocytogenes* infection, 73
Group B streptococcus:
persistence in macrophages, 86

HER-2/neu protein, human:
and tolerance to rat neu, 192
HLA-DM in B-lymphoblastoid cells:
trafficking, 505

Immunoglobulin E antibody:
inhibition by $CD8^+$ T cells, 230
Immunoglobulin M heavy chain constant region gene, bovine:
characterization, 581
Inflammation, carrageenan induced:
role of poly ADP-ribose synthetase, 96
Inflammatory reactions:
in human malaria, 264
Inhibition of allergic response:
by *Mycobacterium vaccae*, 307
Interferon-ε:
in murine *Staphylococcus aureus* infection, 80
Interleukin-6 and interleukin-6α expansion:
by B-cell hybridomas, 498
Intranasal immunization:
with herpes simplex virus vaccines, 563
ISCOMS:
adjuvant properties in the absence of interleukin-4 and interferon-γ, 556

Lactic dehydrogenase virus:
effects on responses to *Nippostrongylus brasiliensis*, 540
Lymphocyte trafficking:
memory $CD4^+$ T cells, 447

Macrophages:
differential priming by Gram-negative bacteria, 323
Major histocompatibility complex class II isotype expression:
differential regulation, 33
Mannan-binding lectin, chicken:
cloning and sequencing of cDNA, 421
Mast cells, human:
cytokine expression, 289
Mast cells, rat:
phenotypic characterization, 533
Monoclonal antibodies, human:
generated in human/mouse chimeras (trimera), 154
Monoclonal antibodies, human, reactivity:
and charge of heavy-chain V region, 129
Mycobacterium tuberculosis:
genetics of resistance in mice, 270

Nasal-associated lymphoid tissue:
activation by microparticles, 249
Natural killer cells, human:
elevation of cAMP levels, 415
Natural killer cytotoxicity:
enhanced by cathepsin G, 115
NC-2 receptor:
and natural cytotoxicity, 122
Nitric oxide:
and protective immunity to *Schistosoma mansoni*, 55
production by infected human polymorphonuclear leucocytes, 275
Nitric oxide-mediated cytotoxic effects:
independence from β_2 integrin-mediated adhesion, 102
NK1.1 signalling:
apoptosis in natural killer cells and interleukin-4 production in NK T cells, 296

Orf virus OV20.0L gene product:
and interferon resistance, 335

Peritoneal macrophages:
in hu-PBL-severe combined immunodeficiency mice, 524
Protein kinase C:
mediation of interferon-β mRNA expression in cytotoxic T cells, 455

Somatic mutation of human immunoglobulin V_H6 rearrangements, 171
Species-specific peptides:
from *Mycobacterium leprae*, 64
Superantigen-induced suppression:
and interleukin-12, 485
Suppressor macrophages:
in aged mice, 409
Synovial fibroblasts:
targets for staphylococcal enterotoxin-induced T-cell cytotoxicity, 20

T-cell clones:
antigen direction into gingival tissues, 11
T-cell epitope, common:
mycobacterial, 314
T-cell proliferation:
induction by simultaneous cross-linking of CD6 and CD28, 358
T-cell receptor:
Vδ-Jα rearrangements in human thymocytes, 208
T-cell-receptor peptide:
and immunosenescence, 462
T-cell-receptor transcripts:
and distal Vβ promoters, 213
T-cell subsets:
$CD8^+$ granzyme B^+ in healthy individuals, 383
co-stimulation requirements, 366
in vaccine-induced immunity to *Bordetella pertusis*, 1
T-helper cytokine:
and specificity of T-cell response to heat shock protein, 493
Tolerogenicity:
of polyethylene glycol conjugates, 200

$Vβ8^+$ and $γδ^+$ cells:
in contact sensitivity, 469
V_H gene sequence analysis:
of B-cell tumours, 162